

UNIVERSIDADE DE SÃO PAULO
Instituto de Ciências Matemáticas e de Computação
ISSN 0103-2577

**BOURGIN-YANG VERSIONS OF THE BORSUK-ULAM
THEOREM FOR (H;G)-COINCIDENCES**

WACŁAW MARZANTOWICZ
DENISE DE MATTOS
TACIANA OLIVEIRA SOUZA

Nº 427

NOTAS DO ICMC

SÉRIE MATEMÁTICA



São Carlos – SP
Out./2016

BOURGIN-YANG VERSIONS OF THE BORSUK-ULAM THEOREM FOR (H, G)-COINCIDENCES

WACŁAW MARZANTOWICZ, DENISE DE MATTOS AND
TACIANA OLIVEIRA SOUZA

ABSTRACT. Let $G = \mathbb{Z}_{p^k}$ be a cyclic group of prime power order, or repectively $G = \mathbb{Z}_p^k$ be the p -torus of rank k . We estimate the size of the (H, G) -coincidences set of a continuous map from $S(V)$ into a real vector space W' .

1. INTRODUCTION

Let G be a finite group which acts freely on a space X and let $f : X \rightarrow Y$ be a continuous map from X into another space Y . If H is a subgroup of G , then H acts on the right on each orbit Gx of G as follows: if $y \in Gx$ and $y = gx$, $g \in G$, then $hy = gh^{-1}x$. A point $x \in X$ is said to be a (H, G) -coincidence point of f (as introduced by Gonçalves, Pergher and Jaworowski in [3]) if f sends every orbit of the action of H on the G -orbit of x to a single point. Let us denote by $A(f, H, G)$ the set of all (H, G) -coincidence points. Of course, if H is the trivial subgroup, then every point of X is a (H, G) -coincidence. If $H = G$, this is the usual definition of G -coincidence, that is,

$$A(f, G, H) = A(f) = \{x \in X \mid f(x) = f(gx), \text{ for all } g \in G\}.$$

Borsuk-Ulam theorems type consists in estimating the dimension of the set $A(f, H, G)$. For the case that $G = H = \mathbb{Z}_2$, $X = S^n$ and $Y = \mathbb{R}^n$, we have the classical Borsuk-Ulam theorem [2].

Let $G = \mathbb{Z}_{p^k}$ be a cyclic group of prime power order, $k \geq 1$. For given two powers $1 \leq m \leq n \leq p^{k-1}$ of p we set

$$(1) \quad \mathcal{A}_{m,n} := \{G/H \mid H \subset G; m \leq |H| \leq n\},$$

where $|H|$ is the cardinality of H . We shall write \mathcal{A}_X for a set of all the G -orbits of a space X (up to a homeomorphism, thus up to an isomorphism of finite G -sets).

Let V be an orthogonal representation of $G = \mathbb{Z}_{p^k}$, p prime, $k \geq 1$, such that $V^G = \{0\}$, for the set of fixed points of G . For $G = \mathbb{Z}_{p^k}$, with p odd, every nontrivial irreducible orthogonal representation is even dimensional and admits the complex structure ([4]), thus V admit it too. We denote by $d(V) = \dim_{\mathbb{C}} V = \frac{1}{2} \dim_{\mathbb{R}} V$ the integral numerical invariant of V . If $G = \mathbb{Z}_{2^k}$ and V is a real orthogonal representation of G , then we denote $d(V) = \dim_{\mathbb{R}} V$.

Given W' a real vector space and a continuous map $f : S(V) \rightarrow W'$, in this work we estimate the size of $A(f, \mathbb{Z}_{p^i}, \mathbb{Z}_{p^k})$ the \mathbb{Z}_{p^k} -coincidences set of f , as follows.

2010 *Mathematics Subject Classification*. Primary 55M20 Secondary 55M35, 55N91, 57S17.

Key words and phrases. equivariant maps, covering dimension, orthogonal representation, equivariant K -theory.

Theorem 1.1. *Let V be a complex orthogonal representation of the cyclic group $G = \mathbb{Z}_{p^k}$, p prime, $k \geq 1$, such that $V^G = \{0\}$ and let W' be a real vector space. Let $f : S(V) \rightarrow W'$ be a continuous map.*

(1) *If $\mathcal{A}_{S(V)} \subset \mathcal{A}_{1,p^{k-1}}$, then for all $1 \leq i \leq k$,*

$$\dim A(f, \mathbb{Z}_{p^i}, \mathbb{Z}_{p^k}) \geq 2 \left(\frac{d(V) - 1}{p^{k-1}} \right) - (p^k - p^{k-i})d(W').$$

(2) *If $\mathcal{A}_{S(V)} \subset \mathcal{A}_{1,p^{i-1}}$ for some $1 \leq i \leq k$, then*

$$\dim A(f, \mathbb{Z}_{p^i}, \mathbb{Z}_{p^k}) \geq 2 \left(\frac{d(V) - 1}{p^{i-1}} \right) - (p^k - p^{k-i})d(W').$$

Theorem 1.2. *Let V be a real orthogonal representation of the cyclic group $G = \mathbb{Z}_{2^k}$, $k \geq 1$, such that $V^G = \{0\}$ and let W' be a real vector space. Let $f : S(V) \rightarrow W'$ be a continuous map.*

(1) *If $\mathcal{A}_{S(V)} \subset \mathcal{A}_{1,2^{k-1}}$, then for all $1 \leq i \leq k$,*

$$\dim A(f, \mathbb{Z}_{2^i}, \mathbb{Z}_{2^k}) \geq \left(\frac{d(V) - 1}{2^{k-1}} \right) - (2^k - 2^{k-i})d(W').$$

(2) *If $\mathcal{A}_{S(V)} \subset \mathcal{A}_{1,2^{i-1}}$ for some $1 \leq i \leq k$, then*

$$\dim A(f, \mathbb{Z}_{2^i}, \mathbb{Z}_{2^k}) \geq \left(\frac{d(V) - 1}{2^{i-1}} \right) - (2^k - 2^{k-i})d(W').$$

2. BOURGIN-YANG VERSIONS OF THE BORSUK-ULAM THEOREM FOR $G = \mathbb{Z}_{p^k}$

Recently, in [5], the authors proved the following Bourgin-Yang version of the Borsuk-Ulam theorem for complex orthogonal representations of $G = \mathbb{Z}_{p^k}$, p prime, $k \geq 1$

Theorem 2.1. [5, Theorem 3.6] *Let V, W be two complex orthogonal representations of the cyclic group $G = \mathbb{Z}_{p^k}$, p prime, $k \geq 1$, such that $V^G = W^G = \{0\}$. Let $f : S(V) \xrightarrow{G} W$ be an equivariant map and $Z_f := f^{-1}(0) = \{v \in S(V) \mid f(v) = 0\}$. Suppose $\mathcal{A}_{S(V)} \subset \mathcal{A}_{m,n}$ and $\mathcal{A}_{S(W)} \subset \mathcal{A}_{m,n}$. Then*

$$\dim(Z_f) \geq 2 \left(\left(\frac{(d(V) - 1)m}{n} \right) - d(W) \right).$$

They also proved the following Bourgin-Yang version of the Borsuk-Ulam theorem for real orthogonal representations of $G = \mathbb{Z}_{2^k}$, $k \geq 1$.

Theorem 2.2. [5, Theorem 3.9] *Let V, W be two real orthogonal representations of the cyclic group $G = \mathbb{Z}_{2^k}$, $k \geq 1$, such that $V^G = W^G = \{0\}$. Let $f : S(V) \xrightarrow{G} W$ be an equivariant map and $Z_f = f^{-1}(0)$. Suppose that $\mathcal{A}_{S(V)} \subset \mathcal{A}_{m,n}$ and $\mathcal{A}_{S(W)} \subset \mathcal{A}_{m,n}$. Then*

$$\dim(Z_f) \geq \left(\frac{(d(V) - 1)m}{n} \right) - d(W).$$

3. PROOF OF THE MAIN RESULTS

Proof of Theorem 1.1 Let i be fixed, with $1 \leq i \leq k$ and let us consider the real vector space $\bigoplus_{j=1}^{p^k} W'$, which is the direct sum of p^k copies of W' . We have that $\bigoplus_{j=1}^{p^k} W'$ admits an action of the cyclic group $G = \mathbb{Z}_{p^k}$, given by

$$g(w_1, w_2, \dots, w_{p^k}) = (w_2, \dots, w_{p^k}, w_1),$$

for a fixed generator $g \in G$ and for each $(w_1, \dots, w_{p^k}) \in \bigoplus_{j=1}^{p^k} W'$.

Let us denote by $\Delta(W'^{p^{k-i}})$ the diagonal of

$$\bigoplus_{j=1}^{p^k} W' = W'^{p^{k-i}} \oplus \dots \oplus W'^{p^{k-i}}.$$

We have

$$\bigoplus_{j=1}^{p^k} W' = \Delta(W'^{p^{k-i}}) \oplus (\Delta(W'^{p^{k-i}}))^\perp,$$

where $\Delta(W'^{p^{k-i}})^\perp$ is the orthogonal complement of $\Delta(W'^{p^{k-i}})$. Since $\Delta(W'^{p^{k-i}})$ is a $p^{k-i} \dim W'$ - dimensional G -subspace of $\bigoplus_{i=1}^{p^k} W'$, let us observe that $\Delta(W'^{p^{k-i}})^\perp$ is a $(p^k - p^{k-i}) \dim W'$ - dimensional G -subrepresentation of $\bigoplus_{i=1}^{p^k} W'$, for which $(\Delta(W'^{p^{k-i}})^\perp)^G = \{0\}$.

Now, we denote by a_1, \dots, a_r a set of representatives of the left lateral classes of G/\mathbb{Z}_{p^i} , where $r = p^{k-i}$. Consider the map

$$F : S(V) \rightarrow \Delta(W'^{p^{k-i}}) \oplus \Delta(W'^{p^{k-i}})^\perp$$

defined by

$$F(x) = (f(a_1x), \dots, f(a_rx), f(a_1hx), \dots, f(a_rhx), \dots, f(a_1h^{p^i-1}x), \dots, f(a_rh^{p^i-1}x)),$$

for a fixed generator $h \in \mathbb{Z}_{p^i}$. The linear orthogonal projection along the diagonal $\Delta(W'^{p^{k-i}})$ defines a G -equivariant map $\rho : \Delta(W'^{p^{k-i}}) \oplus \Delta(W'^{p^{k-i}})^\perp \rightarrow \Delta(W'^{p^{k-i}})^\perp$. Let us denote by l the composition

$$S(V) \xrightarrow{F} \Delta(W'^{p^{k-i}}) \oplus \Delta(W'^{p^{k-i}})^\perp \xrightarrow{\rho} \Delta(W'^{p^{k-i}})^\perp,$$

with $Z_l = l^{-1}(0) = (\rho \circ F)^{-1}(0) = F^{-1}(\Delta(W'^{p^{k-i}})) = A(f, \mathbb{Z}_{p^i}, \mathbb{Z}_{p^k})$.

For a fixed generator $g \in G$, we can consider

$$h = g^{p^{k-i}} \quad \text{and} \quad a_1 = e, a_2 = g, \dots, a_r = g^{p^{k-i}-1},$$

then F is a G -equivariant map. Moreover

$$(2) \quad \mathcal{A}_{S(\Delta(W'^{p^{k-i}})^\perp)} \subset \mathcal{A}_{1, p^{i-1}} \subset \mathcal{A}_{1, p^{k-1}}.$$

To check the validity of inclusion $\mathcal{A}_{S(\Delta(W'^{p^{k-i}})^\perp)} \subset \mathcal{A}_{1, p^{i-1}}$, it suffices to prove that the cardinality of the orbit $\mathbb{Z}_{p^k} w$ belongs to the set $\{p^k, p^{k-1}, \dots, p^{k-i+1}\}$, for all $w = (w_1, \dots, w_{p^k}) \in$

$S(\Delta(W'^{p^{k-i}})^\perp)$. According to [1, Chapter 1, Proposition 4.1], the cardinality of the orbit $\mathbb{Z}_{p^k}w$ belongs to the set $\{p^k, p^{k-1}, \dots, p, p^0 = 1\}$. Let $w = (w_1, \dots, w_{p^k})$ an element in $S(\Delta(W'^{p^{k-i}})^\perp)$ and suppose that

$$|\mathbb{Z}_{p^k}w| \in \{p^{k-i}, p^{k-i-1}, \dots, p^0 = 1\},$$

that is $|\mathbb{Z}_{p^k}w| = p^j$, for some $0 \leq j \leq k - i$.

Assertion. We have $\mathbb{Z}_{p^k}w = \{w, gw, \dots, g^{p^j-1}w\}$, for a fixed generator g of \mathbb{Z}_{p^k} .

In fact, consider a cyclic group G , $g \in G$ a fixed generator and $\{w, gw, \dots, g^{s-1}w\}$ the maximum set of the s first elements of the orbit Gw that are distinct from each other. By definition of the set $\{w, gw, \dots, g^{s-1}w\}$ we have

$$g^s w \in \{w, gw, \dots, g^{s-1}w\}.$$

Suppose that

$$g^s w = g^i w, \text{ for some } 1 \leq i \leq s - 1,$$

then

$$g^{s-i} w = w, \quad 1 \leq s - i \leq s - 1,$$

but this contradicts the definition of $\{w, gw, \dots, g^{s-1}w\}$.

Now, if $g^t w \in Gw$, for some $t \in \mathbb{N}$, we have $t = ns + r$ with $0 \leq r \leq s - 1$. Therefore

$$g^t w = g^{ns+r} w = g^r (g^{ns} w) = g^r w \in \{w, gw, \dots, g^{s-1}w\},$$

since

$$g^{ns} w = (g^s \cdots g^s) w = w$$

and $0 \leq r \leq s - 1$.

Thus, for a fixed generator g of \mathbb{Z}_{p^k} , we have

$$\begin{aligned} w = g^{p^j} w &= g^{p^j} (w_1, \dots, w_{p^j}, \dots, w_{(p^{k-j-1})p^j+1}, \dots, w_{p^k}) \\ &= (w_{p^j+1}, \dots, w_{2p^j}, \dots, w_{(p^{k-j-1})p^j+1}, \dots, w_{p^k}, w_1, \dots, w_{p^j}) \end{aligned}$$

therefore $w \in \Delta(W'^{p^j})$. Since

$$\Delta(W') \subset \Delta(W'^p) \subset \dots \subset \Delta(W'^{p^{k-i-1}}) \subset \Delta(W'^{p^{k-i}})$$

and $j \in \{0, 1, \dots, k - i\}$, we conclude that

$$w \in \Delta(W'^{p^j}) \subset \Delta(W'^{p^{k-i}}),$$

which is a contradiction because

$$\Delta(W'^{p^{k-i}}) \cap S(\Delta(W'^{p^{k-i}})^\perp) = \emptyset.$$

Thus the Theorem 2.1 implies the claim. \square

Proof of Theorem 1.2 For $G = \mathbb{Z}_{2^k}$, $k \geq 1$, using the same steps of the proof of Theorem 1.1 and applying Theorem 2.2 we have the result. \square

Remark 3.1. *We emphasize that in the Theorems 1.1 and 1.2 the action of G on $S(V)$ is not necessarily free. Moreover, we have an estimate for the size of the set of (H, G) -coincidence points for all subgroups $H = \mathbb{Z}_{p^i}$ of $G = \mathbb{Z}_{p^k}$. These two characteristics make the Theorems 1.1 and 1.2 different of classical results about (H, G) -coincidences.*

4. CASE OF G BEING A p -TORUS

Now, let V, W be two orthogonal representations of the p -torus group $G = \mathbb{Z}_p^k$ of rank $k \geq 1$, p prime, such that $V^G = W^G = \{0\}$ for the sets of fixed points of G . Let $f : S(V) \rightarrow W'$ be a G -equivariant map, in [6, Theorem 2.7] we have the following estimate for the covering dimension of Z_f

$$\dim(Z_f) \geq d(V) - d(W) - 1.$$

Using this result we can estimate the covering dimension of the set of (H, \mathbb{Z}_p^k) -coincidences, as follows.

Corollary 4.1. *Let V, W' be two orthogonal representation of the group $G = \mathbb{Z}_p^k$, such that $V^G = W'^G = \{0\}$. Let $f : S(V) \rightarrow W'$ be a G -equivariant map, then*

$$\dim A(\mathbb{Z}_p^i, \mathbb{Z}_p^k) \geq d(V) - (p^k - p^{k-i})d(W') - 1$$

for all $1 \leq i \leq k$.

Proof: Let i be fixed, with $1 \leq i \leq k$. As in the proof of the Theorem 1.1, we consider a map

$$F : S(V) \rightarrow \bigoplus_{j=1}^{p^k} W'$$

defined by

$$F(x) = (f(a_1 h_1 x), \dots, f(a_r h_1 x), f(a_1 h_2 x), \dots, f(a_r h_2 x), \dots, f(a_1 h_{p^i} x), \dots, f(a_r h_{p^i} x)),$$

where a_1, \dots, a_r are representatives of the left lateral classes of G/\mathbb{Z}_p^i , $r = p^{k-i}$ and $\mathbb{Z}_p^i = \{h_1, \dots, h_{p^i}\}$.

Considering the diagonal action of G on $\bigoplus_{j=1}^{p^k} W'$, F becomes G -equivariant. Thus the composition

$$S(V) \xrightarrow{F} \Delta(W'^{p^{k-i}}) \oplus \Delta(W'^{p^{k-i}})^\perp \xrightarrow{\rho} \Delta(W'^{p^{k-i}})^\perp,$$

is also G -equivariant. Finally, we apply [6, Theorem 2.7] to estimate the size of the set $Z_{\rho \circ F} = A(\mathbb{Z}_p^i, \mathbb{Z}_p^k)$. \square

REFERENCES

- [1] Bredon, G., *Introduction to compact transformation groups*, Vol. **46**, Academic Press, New York and London, (1972).
- [2] Borsuk, K., *Drei Sätze über die n -dimensionale euklidische Sphäre*, Fund. Math. **20** 177–190, 1933.
- [3] Gonçalves, D.L.; Jaworowski, J.; Pergher, Pedro L.Q., *Measuring the size of the coincidence set*, Topology Appl. **125** (2002), 465–470.

- [4] Serre, J. P. *Linear representations of finite groups*. Translated from the second French edition by Leonard L. Scott., Graduate Texts in Mathematics 42, Springer-Verlag, New York-Heidelberg, (1977).
- [5] Waclaw Marzantowicz, D. de Mattos, Edivaldo L. dos Santos, *Bourgin-Yang version of the Borsuk-Ulam theorem for \mathbb{Z}_p^k -equivariant maps*, Algebraic & Geometric Topology, to appear 2012.
- [6] Waclaw Marzantowicz, D. de Mattos, Edivaldo L. dos Santos, *Bourgin-Yang versions of the Borsuk-Ulam theorem for p -toral groups*.

FACULTY OF MATHEMATICS AND COMPUTER SCI., ADAM MICKIEWICZ UNIVERSITY OF POZNAŃ, UL. UMULTOWSKA 87, 61-614 POZNAŃ, POLAND.

E-mail address: marzan@amu.edu.pl

INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO, DEPARTAMENTO DE MATEMÁTICA, CAIXA POSTAL 668, SÃO CARLOS-SP, BRAZIL, 13560-970.

E-mail address: deniseml@icmc.usp.br

E-mail address: tacioli@icmc.usp.br